

the size of BLRs of low luminous Active Galactic Nuclei

Xue-Guang, Zhang^{1,2*}, Dultzin-Hacyan, Deborah.¹, Ting-Gui, Wang²

¹*Instituto de Astronomía, Universidad Nacional Autónoma de México, Apdo Postal 70-264, México D. F. 04510, Mexico*

²*Center for Astrophysics, Department of astronomy and Applied Physics, University of Science and Technology of China, Hefei, Anhui, P.R.China*

ABSTRACT

We study the size of BLRs of low luminosity Active Galactic Nuclei, also called ‘dwarf AGN’, defined as having ($L_{H\alpha} \leq 10^{41} \text{ erg} \cdot \text{s}^{-1}$). We more than double the sample size analyzed previously (Wang & Zhang 2003, hereafter Paper I). In this study we first confirm our previous result that the sizes of BLRs of low luminosity AGN are larger than the ones expected from the empirical relation $R_{BLRs} - L_{H\alpha}$ valid for ‘normal’ AGN: Seyfert 1s and quasars, except for the objects with accretion rate $\dot{m}_{H\alpha} > 10^{-5.5}$. Second, we find a positive correlation between the line width of the narrow emission line (as tracer of velocity dispersion and thus bulge and black hole mass) and the size of BLRs for both normal and low luminosity AGN. In this paper we find a non-linear dependence of the BLRs sizes of low luminosity AGN on BH masses. We also show that their sizes of BLRs are more strongly dominated by the ‘specific accretion rate’ $\dot{m}_{H\alpha}$ defined as $\dot{m}_{H\alpha} = L_{H\alpha}/L_{Edd}$, than by the masses of their central black holes. As an expected result, the distance of emission regions of low-ionization broad $H\alpha$ of NGC 4395 should be consistent with the value from the empirical relation of $R_{BLRs} - L_{H\alpha}$, according to the high accretion rate

Key words: Galaxies:Active – Galaxies:nuclei – Galaxies:low luminosity

1 INTRODUCTION

Broad emission lines are one of the prominent properties of type 1 Active Galactic Nuclei (AGN). However, due to the limit of the resolution of present, and even foreseeable astronomical instruments, we have no way to resolve the geometric structure of broad emission line regions (BLRs) by direct observations. Throughout this paper we shall use the term ‘size of BLRs’ as equivalent of distance of the BLRs to the central source. As the result of efforts of hundreds of astronomers during more than ten years, the sizes of the BLRs have been estimated for more than 30 AGN by the Reverberation Mapping technique (Blandford & McKee 1982; Peterson 1993; Kaspi et al. 2005). Using a sample of 35 objects, 15 PG quasars and 21 Seyfert 1 galaxies, a strong correlation between the size of BLRs and the continuum luminosity has been found (Kaspi et al. 2000; 2005). The empirical relation $R_{BLRs} - L_{5100\text{\AA}}^\alpha$ indicates that both the ionization continuum and the electron density of the BLRs for normal AGN are not much different, and provides a convenient way to estimate the central BH masses for AGN under the assumption of Virialization (Greene & Ho 2005b; Ovcharov et al. 2005; Wu et al. 2004; McLure & Jarvis 2004; 2002; Marziani et al. 2003), especially for objects with low and intermediate

redshifts and high S/N spectra (Sulentic et al. 2006). Due to the difficulty to measure the stellar velocity dispersion, the relation $M_{BH} - \sigma$ which can be extrapolated for intermediate BH masses in the range of $10^4 - 10^6 M_\odot$ (Barth et al. 2005; Greene & Ho 2004), cannot be applied to higher luminosity and higher redshift AGN (Tremaine et al. 2002; Ferrarese & Merritt 2001; Gebhardt, Bender et al. 2000).

In Paper I we have shown that the empirical relation $R_{BLRs} - L_{H\alpha}^\alpha$ is not valid for a special kind of AGN: low luminosity active galaxies with $L_{H\alpha} < 10^{41} \text{ erg} \cdot \text{s}^{-1}$ selected from Ho et al. (1997a, 1997b), and called ‘dwarf AGN’ by them. The deviation of this empirical relation for low luminosity AGN may be explained in terms of a lower ionization parameter and/or lower electron density of BLRs in these type of AGN (although see discussion paper I). Broad emission lines (often only $H\alpha$) can be detected in the spectra of these type of galaxies indicating the presence of activity in their nuclei, and there is also other evidence for the presence of massive black holes in low luminosity AGN (Nakai et al. 1993; Sarzi et al. 2001). In the case of low luminosity AGN with intermediate BH masses $M_{BH} \sim 10^4 - 10^6 M_\odot$, however, several objects have been studied individually in order to check if they follow the correlation $M_{BH} - \sigma$: POX 52 (Barth et al. 2004), NGC 4395 (Kraemer et al. 1999; Filippenko & Ho 2003), NGC4051 (Shemmer et al. 2003) and various objects selected from SDSS (Greene & Ho 2004). The consistency between the BH masses estimated using

* xguang@astroscu.unam.mx

the relation $M_{BH} - \sigma$ and estimated under the assumption of virialization, indicates that the size of BLRs for these AGN with intermediate BH masses must obey the empirical relation $R_{BLRs} - L_{5100\text{\AA}}$. Thus the question of whether all the sizes of BLRs of low luminosity AGN obey the empirical relation is an interesting one. In this paper, we use a larger sample of low luminosity AGN to further investigate this question, in order to find a more consistent explanation for the deviation of the empirical relation for low luminosity AGN reported in Paper I. The next section presents the sample. The results are shown in Section III. Section IV gives the discussion and conclusion. In this paper, the cosmological parameters $H_0 = 70 \text{ km} \cdot \text{s}^{-1} \text{ Mpc}^{-1}$, $\Omega_\Lambda = 0.7$ and $\Omega_m = 0.3$ have been adopted.

2 SAMPLE

In a spectroscopic survey of more than four hundred nearby galaxies, there are 34 AGN, of which the broad emission lines have been detected with confidence, and 12 AGN with less certainty (Ho et al. 1997a, 1997b). In paper I, we selected 22 low luminosity AGN with broad emission $H\alpha$, 4 Seyfert galaxies and also 18 LINERs selected from the sample of Ho et al. (1997a, 1997b), to study the size of BLRs.

As in paper I, we select low luminosity AGN according to the criterion that $L_{H\alpha} \leq 10^{41} \text{ erg} \cdot \text{s}^{-1}$ (see also Ho et al. 1997a; 1997b). We first select 42 low luminosity AGN from Ho et al. (1997b), including the objects NGC4051 and NGC4395, for which the Galactic and internal reddening have been corrected using the values given by Ho et al. (1997a): The internal color excess can be found in their Table 4, the Galactic color excess can be found in their Table 11. Although there are 46 objects for which broad $H\alpha$ lines have been detected, there are four objects NGC1275, NGC4151, NGC3516 and NGC5548 with a luminosity of $H\alpha$ larger than $10^{41} \text{ erg} \cdot \text{s}^{-1}$, thus, we reject them for our sample. Furthermore, in paper I, there are 7 objects with double-peaked broad low-ionization emission lines (dbp emitters), we excluded these objects from our new sample. Finally, we also selected low luminosity AGN from the literature according to the above defined criterion for $L_{H\alpha}$. We find four additional objects which can be selected as candidate, POX 52 (Barth et al. 2004), SDSS J024912.86-081525.6, SDSS J143450.62+033842.5 and SDSS J233837.10-002810.3 (Greene & Ho 2004, 2005b). In order to estimate the BH masses, we compile the velocity dispersion of the bulge for each object. If there are no data for the velocity dispersion of the bulge, we use the line width of the strong narrow emission line $[\text{NII}]\lambda 6583\text{\AA}$. The strong correlation between the line width of the narrow emission lines and the stellar velocity dispersion of the bulge has been studied by several authors (e.g. Nelson & Whittle 1996; Greene & Ho 2005a). The sample objects are listed in Table 1.

Finally, we select the 35 objects with accurate determinations for the sizes of their BLRs estimated from reverberation mapping technique, in order to compare the difference for the two kinds of AGN, low luminosity and normal AGN. The 35 normal AGN can be found in Kaspi et al. (2005) and Peterson et al. (2004). We do not list them again.

3 RESULTS

3.1 The size of BLRs of dwarf AGN

As described in paper I, we estimate the size of BLRs of low luminosity AGN using:

$$M_{BH} = 10^{8.13 \pm 0.06} \left(\frac{\sigma}{200 \text{ km} \cdot \text{s}^{-1}} \right)^{4.02 \pm 0.32} M_\odot \quad (1a)$$

$$R = q \times \frac{G \times M_{BH}}{FWHM_{H\alpha B}^2} \quad (1b)$$

where $FWHM_{H\alpha B}$ is the line width of broad $H\alpha$, and q is a factor related to the orbits of the emission clouds. It is also pointed out that $q = \frac{4}{3}$ is the most convenient value discussed in the literature. In more recent literature, however, Onken et al. (2004) found that $M_{BH} = 5.5 \times \frac{R\sigma_{line}^2}{G}$ and Peterson et al. (2004) found the mean value of $FWHM/\sigma = 2.03 \pm 0.59$. Thus, here we take $q = \frac{1}{5.5/2.03^2} = 0.75$ rather than $q = \frac{4}{3}$.

There are 38 low luminosity AGN, for which the stellar velocity dispersions can be compiled from the recent literature. There is one object, NGC4395, of which the upper limit value of stellar velocity dispersion, $\sigma < 30 \text{ km} \cdot \text{s}^{-1}$, can be selected from Fillipenko & Ho (2003). In order to estimate the BH masses for the other seven, we first check the correlation between the stellar velocity dispersion and the line width of narrow emission lines. We adopt the $[\text{NII}]\lambda 6583\text{\AA}$ measurements of Ho et al. (1997a). We show this correlation in Fig 1. The Spearman Rank correlation coefficient is 0.77 with $P_{null} \leq 8.8 \times 10^{-8}$. The mean value of $FWHM/\sigma$ is about 2.13 ± 0.12 . Here, we do not consider the object NGC 1068, because of the special geometric structure of its narrow emission line region (NLR), which has been observed with HST. The NLR of NGC 1068 consists of a large set of clouds (Macchetto et al. 1994; Capetti et al. 1995; Dietrich & Wagner 1998; Cecil et al. 2002) which leads to a larger than mean value of line width for narrow emission lines. The strong correlation seen in Fig 1 indicates that we can confidently estimate the BH masses for the eight low luminosity AGN without velocity dispersions using the line width of $[\text{NII}]\lambda 6583\text{\AA}$. Somewhat larger stellar velocity dispersions than those derived from $\sigma_{[\text{NII}]}$ are perhaps due to a larger distance of the $[\text{NII}]$ emission line clouds to the central source. Thus, the sizes of BLRs for these eight low luminosity AGN can also be estimated. The sizes of the BLRs are listed in Table 1. Notice the case of the low luminosity AGN NGC 4395 (because the size of BLRs has been measured by Reverberation Mapping technique by Peterson et al. (2005), thus we list the object individually) for which the BH mass estimated from the upper limit of stellar velocity dispersion is about $6.6 \pm 3.1 \times 10^4 M_\odot$ (we think the upper limit of stellar velocity from Keck high-resolution spectra is more reliable than the upper limit of line width of $[\text{NII}]$), which is about four to six times smaller than the BH mass $3.6 \pm 1.1 \times 10^5 M_\odot$ derived recently under the assumption of Virialization and using reverberation mapping by Peterson et al. (2005).

We can now compare the empirical relation $R_{BLRs} - L_{H\alpha}$ for low luminosity AGN to that of normal AGN. As we can see from Figure 2, the result from paper I is confirmed. The size of BLRs of low luminosity AGN is larger than the prediction of the empirical relation $R_{BLRs} - L_{H\alpha}$. The mean value of $\xi = \log\left(\frac{L_{H\alpha}/10^{44} \text{ erg} \cdot \text{s}^{-1}}{(R_{BLRs}/100 \text{ light-days})^2}\right)$ is about -4.19 ± 0.23

for low luminosity AGN except NGC4395. The Spearman Rank correlation coefficient is 0.06 with $P_{null} \sim 0.72$. For NGC 4395, the object mentioned above, the size of the BLRs is about 0.04 light-days derived directly from the variation of the CIV emission line, which is much smaller than the upper limit value derived from the upper limit BH masses using broad H α : 1.28 light-days. However, the emission regions of high-ionization CIV should be nearer to the central black hole than the regions of low-ionization H α . We do notice, however, that there are some low luminosity AGN including four objects with intermediate BH masses, for which the sizes of the BLRs do obey the empirical relation $R_{BLRs} - L_{H\alpha}$. Thus, it is important to further investigate which parameter dominates the size of BLRs of both low luminosity and normal AGN.

3.2 BH masses and accretion rates

Due to the strong correlation between BH mass and the continuum luminosity, there is also a strong correlation between BH mass and the size of BLRs for normal AGN. We investigate if the same holds for low luminosity AGN, which can be somehow expected given that the sizes of the BLRs are also calculated from BH masses.

The correlations between M_{BH} and $L_{H\alpha}$ are shown in Figure 3. The Spearman Rank correlation coefficient is -0.09 with $P_{null} \geq 51\%$ for low luminosity AGN except NGC4395, and it is 0.87 with $P_{null} \sim 1.15 \times 10^{-11}$ for normal AGN. Thus, we see that the correlations are different. Next, we show the correlations between M_{BH} and R_{BLRs} in Figure 4. The Spearman Rank correlation coefficient is 0.94 with $P_{null} \sim 4.66 \times 10^{-22}$ for low luminosity AGN except NGC4395, and 0.90 with $P_{null} \sim 1.12 \times 10^{-13}$ for the 35 normal AGN. The unweight linear relations for low luminosity AGN except NGC4395 and normal AGN are:

$$\log(R_{BLRs})(low) = -4.31 \pm 1.14 + 0.78 \pm 0.15 \log(M_{BH}) \quad (2a)$$

$$\log(R_{BLRs})(normal) = -4.36 \pm 1.98 + 0.73 \pm 0.25 \log(M_{BH}) \quad (2b)$$

A correlation between M_{BH} and R_{BLRs} can be expected, because of the fundamental formula $M_{BH} = q \frac{R_{BLRs} FWHM_{H\alpha}^2}{G}$. However, the results above show that there is no simple linear relation $R_{BLRs} \propto M_{BH}^1$, instead $R_{BLRs} \propto M_{BH}^\alpha$ with $\alpha < 1$. Thus we can still say that there is a correlation between BH masses and line width.

We shall next define and calculate a parameter related to the dimensionless accretion rate, $\dot{m} = L_{bol}/L_{Edd}$, where $L_{bol} \sim 9 \times L_{5100\text{\AA}}$ is the bolometric luminosity for Normal AGN (Wandel et al. 1999; Kaspi et al. 2000) and L_{Edd} is the Eddington luminosity $L_{Edd} = 1.38 \times 10^{38} M_{BH}/M_\odot \text{ erg} \cdot \text{s}^{-1}$. It is difficult to determine accurately the continuum luminosity of the nucleus for low luminosity AGN, because of the dominant component from stellar light and the lack of big blue bump.

Unfortunately, it was also shown by Ho (1999) that for low luminosity AGN, the bolometric luminosity cannot be estimated as $9 \times L_{5100\text{\AA}}$, since for example the ratio of bolometric luminosity to continuum luminosity at 5100Å for the low luminosity AGN M81 (NGC 3031) is about 10, whereas the same ratio for another low luminosity AGN,

NGC 4261, is more than 500!. Greene & Ho (2005b) found a strong correlation between the optical continuum luminosity and the luminosity of H α (including both broad and narrow components) using a sample of normal AGN selected from SDSS, but we cannot be sure that the same correlation applies to low luminosity AGN. In what follows, we use the total (broad + narrow) luminosity of H α to define and calculate another dimensionless accretion rate: $\dot{m}_{H\alpha} = L_{H\alpha}/L_{Edd}$.

The correlation between the size of BLRs and the dimensionless accretion rate $\dot{m}_{H\alpha}$ is shown in Figure 5. The Spearman Rank correlation coefficient is -0.71 with $P_{null} \leq 3.48 \times 10^{-8}$ for low luminosity AGN except NGC4395. For the normal 35 AGN, we select the 12 objects with measured stellar velocity dispersions to calculate the dimensionless accretion rate $\dot{m}_{H\alpha} = L_{H\alpha}/L_{Edd}$, and the luminosities of H α for the 12 objects are estimated from the continuum luminosity by the correlation found by Greene & Ho (2005b):

$$L_{H\alpha} = 5.25 \pm 0.02 \times 10^{42} \left(\frac{L_{5100\text{\AA}}}{10^{44} \text{ erg} \cdot \text{s}^{-1}} \right)^{1.157 \pm 0.005} \text{ erg} \cdot \text{s}^{-1} \quad (3)$$

The correlation coefficient is 0.76 with $P_{null} \leq 0.004$ for the 12 normal AGN. The unweighted linear relation for low luminosity AGN is:

$$\log(R_{BLRs}) = -1.37 \pm 0.73 - 0.47 \pm 0.11 \log(L_{H\alpha}/L_{Edd}) \quad (4)$$

Due to the small number of normal AGN, we cannot confirm whether the different correlation between R_{BLRs} and $\dot{m}_{H\alpha}$ for low luminosity and normal AGN is really true. However, we can confirm that the few low luminosity AGN for which the sizes of the BLRs are consistent with the empirical correlation $R_{BLRs} - L_{H\alpha}$, have large values of $\dot{m}_{H\alpha}$ than $\sim 10^{-5.5}$. These low luminosity AGN have been marked with names or shown as triangles in Figure 2.

4 DISCUSSION AND CONCLUSIONS

BH masses estimated from stellar velocity dispersion are reliable for low luminosity AGN. If the size of the BLRs of low luminosity AGN with BH masses larger than $10^6 M_\odot$ would obey the correlation $R_{BLRs} - L_{con}^\alpha$, their lower continuum luminosity would lead to at least one order of magnitude larger width of broad lines than what is observed in our sample. Such broader emission lines cannot be detected in high resolution spectra of HST. With respect to the issue of the detectability limit of broad emission lines for ground based observations, we wish to point out as an example that Barth et al. (2001) observed the BLRs for one low luminosity LINER: NGC 4579, with the STIS on HST, and detected broad H α with the same width, $FWHM = 2300 \text{ km} \cdot \text{s}^{-1}$, as the value detected by Ho et al. (1997b). In the following section, we will see that the line widths of broad H α in our sample selected from Ho et al. (1997b) are not much less than $2300 \text{ km} \cdot \text{s}^{-1}$, thus we think these line widths are not limited by the Palomar spectra detection capability. We cannot rule out however, the possibility that there may be two distinct BLRs for at least some low luminosity AGN: The one observed in the reported objects, and a very broad one near to the black hole, possibly emitting double peaked lines, which is presently undetectable.

In this paper we have investigated the role of luminosity in the structure of the BLR. In Paper I it was found that low-luminosity AGN have abnormally large BLRs for a given luminosity, compared to more luminous 'normal' AGN. The current paper expands the sample of low-luminosity AGN to include not only massive BHs in low accretion states, but also intermediate-mass black holes radiating close to their Eddington limits. The latter, with possibly one exception, despite their comparably low luminosities, appear to have 'normal-sized' BLRs. The dependence of BLRs structure on luminosity and Eddington ratio may have important implications for BH mass measurements and AGN physics in general. According to the high accretion rate of $\dot{m}_{H\alpha}$ of NGC4395, we can expect that the size of BLRs, particularly the emission regions of low-ionization broad $H\alpha$ should be consistent with the value from the empirical relation of $R_{BLRs} - L_{H\alpha}$.

ACKNOWLEDGEMENTS

ZXG gratefully acknowledges the postdoctoral scholarships offered by la Universidad Nacional Autonoma de Mexico (UNAM). D. D-H acknowledges support from grant IN100703 from DGAPA, UNAM. This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. This research has also made use of the HyperLeda.

REFERENCES

- Arribas S., Mediavilla E., Garcia-Lorenzo B., Del Burgo C., 1997, *ApJ*, 490, 227
- Barth A. J., Ho L. C., Filippenko A. V. & Rix H. W., 2001, *ApJ*, 546, 205
- Barth A. J., Ho L. C. & Sargent W. L. W., 2002, *AJ*, 124, 2607
- Barth A. J., Ho L. C., Rutledge R. E. & Sargent W. L. W., 2004, *ApJ*, 607, 90
- Barth A. J., Greene J. E. & Ho L. C., 2005, *ApJ*, 619, L151
- Bernardi M., Alonso M. V., da Costa L. N., Willmer C. N. A., Wegner G., Pellegrini P. S., et al., 2002, *AJ*, 123, 2990
- Blandford R. D. & McKee C. F., 1982, *ApJ*, 255, 419
- Capetti A., Axon D. J., Macchetto F. D., Sparks W. B. & Boksenberg A., 1995, *APJ*, 446, 155
- Cecil G., Dopita M. A., Groves B. et al., 2002, *APJ*, 568, 627
- DeSouza R. E., Barbuy B. & dos Anjos S., 1993, *AJ*, 105, 1737
- Dietrich M. & Wagner S. J., 1998, *A&A*, 338, 405
- Ferrarese L. & Merritt D., 2001, *MNRAS*, 320, L30
- Filippenko A. V. & Ho L. C., 2003, *ApJ*, 588, L13
- Fisher D., 1997, *AJ*, 113, 950
- Garcia-Lorenzo B., Mediavilla E., Arribas S. & Del Burgo C., 1997, *ApJ*, 483, L99
- Gebhardt K., Bender R., Bower G., et al., 2000, *ApJ*, 439, L13
- Greene J. E. & Ho L. C., 2004, *ApJ*, 610, 722
- Greene J. E. & Ho L. C., 2005a, *ApJ*, 627, 721
- Greene J. E. & Ho L. C., 2005b, *ApJ*, 630, 122
- Heraudeau P. H. & Simien F. 1998, *A&AS*, 133, 317
- Ho L. C., Filippenko A. & Sargent W. L. W., 1997a, *ApJS*, 112, 315
- Ho L. C., Filippenko A. & Sargent W. L. W., 1997b, *ApJS*, 112, 391
- Ho L. C., 1999, *ApJ*, 516, 672
- Jore K.P., Broeils A.H. & Haynes M.P., 1996, *AJ*, 112, 438
- Kaspi S., Smith P. S., Netzer H., et al., 2000, *ApJ*, 533, 631
- Kaspi S., Maoz D., Netzer H., Peterson B. M., et al., 2005, *ApJ*, 629, 61
- Kraemer S. B., Ho L. C., Crenshaw D. M., et al., 1999, *ApJ*, 520, 564
- Lewis K. T. & Eracleous M., 2006, *astro-ph/0601398*
- Macchetto F. D., Capetti A., Sparks W. B., Axon D. J. & Boksenberg A., 1994, *APJ*, 435, L15
- Marziani P., Zamanov R. K., Sulentic J. W. & Calvani M., 2003, *MNRAS*, 345, 1133
- McElory D. B., 1995, *ApJS*, 100, 105
- McLure R. J. & Jarvis M. J., 2002, *MNRAS*, 337, 109
- McLure R. J. & Jarvis M. J., 2004, *MNRAS*, 353, 45
- Nakai N., Inoue M. & Miyoshi M., 1993, *Nature*, 361, 45
- Nelson C. H. & Whittle M., 1995, *ApJS*, 99, 67
- Nelson C. H. & Whittle M., 1996, *APJ*, 465, 96
- Oliva E., Origlia L., Maiolino R. & Moorwood A. F. M., 1999, *A&A*, 350, 9
- Onken C. A., Ferrarese L., Merritt D., Peterson B. M., et al., 2004, *ApJ*, 615, 645
- Ore C. D., Faber S. M., Gonzalez J., Stoughton R. & Burstein D., 1991, *ApJ*, 366, 38
- Ovcharov E., Ivanov V. D., Nedialkov P., 2005, *gbha.conf*, 1340
- Peterson B. M., 1993, *PASP*, 105, 247
- Peterson B. M., Ferrarese L., Gilbert K. M., Kaspi S., et al., 2004, *ApJ*, 613, 682
- Peterson B. M., Bentz M. C., Desroches L. B., et al., 2005, *ApJ*, 632, 799
- Proctor R. N. & Sansom A. E., 2002, *MNRAS*, 333, 517
- Prugniel P., Maubon G. & Simien F., 2001, *A&A*, 366, 68
- Sarzi M., Rix H., Shields J. C., et al., 2001, *ApJ*, 550, 65
- Schechter P. L. 1983, *ApJS*, 52, 425
- Shemmer O., Uttley P., Netzer H. & Mchaardy I. M., 2003, *MNRAS*, 343, 1341
- Simien F. & Prugniel P., 1997, *A&AS*, 126, 15
- Simien F. & Prugniel P., 2002, *A&A*, 384, 371
- Smith E. P., Heckman T. M. & Illingworth G. D., 1990, *ApJ*, 356, 399
- Sulentic J. W., Repetto P., Stirpe G. M., Marziani P., Dultzin-Hacyan D. & Calvani M., 2006, *A&A*, in press
- Terlevich E., Diaz A. I. & Terlevich R. 1990, *MNRAS* 242, 271
- Trager S. C., Worthey G., Faber S. M., et al., 1998, *ApJS*, 116, 1
- Tremaine S., Gebhardt K., Bender R., Bower G., et al., 2002, *ApJ*, 574, 740
- Vega Beltran J. C., Pizzella A., Corsini E. M., Funes J. G., Zeilinger W. W., Beckman J. E. & Bertola F., 2001, *A&A*, 374, 394
- Wandel A., Peterson B. M. & Malkan M. A., 1999, *ApJ*, 526, 579
- Wang T. G. & Zhang X. G., 2003, *MNRAS*, 340, 793
- Wu X. B., Wang R., Kong M. Z., et al., 2004, *A&A*, 424, 793
- de Zeeuw P. T., et al., 2002, *MNRAS*, 329, 513

Table 1. Data of Sample

Name	Dis Mpc	$\log(H\alpha)$ erg · s ⁻¹	σ km · s ⁻¹	FWHM(H α) km · s ⁻¹	FWHM([NII]) km · s ⁻¹	$\log(R_{BLR})$ light-day	Ref.
NGC315	65.8	39.87	310±3	2000	402	2.87	T98
NGC1052	17.8	40.25	215±7	1950	482	2.19	BHS02
NGC1161	25.9	39.38	288	3000	518	2.45	M95
NGC2681	13.3	39.55	111±16	1550	265	1.31	M95, OF91
NGC2787	13.0	38.76	202±5	2050	382	2.11	BHS02
NGC2985	22.4	38.79	185±10	2050	263	1.95	S01
NGC3031	1.4	39.17	173±5	2650	210	1.61	Ve01
NGC3226	23.4	39.09	203±4	2000	465	2.14	PS02
NGC3642	27.5	39.46	109±11	1250	167	1.46	TDT
NGC3718	17.0	37.45	178±19	2350	371	1.77	HS98
NGC3998	21.6	40.57	297±5	2150	401	2.74	F97
NGC4036	24.6	39.17	166±5	1850	359	1.85	F97
NGC4143	17.0	39.13	211±6	2100	307	2.16	SP02
NGC4203	9.7	38.59	167±3	1500	346	2.05	BHS02
NGC4258	6.8	38.88	134	1700	296	1.29	p01
NGC4278	9.7	38.98	251±13	1950	479	2.53	BHS02
NGC4450	16.8	38.83	80	2300	301	0.39	Z02
NGC4452	2.4	38.15	268±12	3700		2.09	SP97
NGC4579	16.8	39.45	165±4	2300	376	1.65	BHS02
NGC4636	17.0	38.36	208±8	2450	333	2.01	BA02
NGC5077	40.6	39.23	275±39	2300	491	2.55	T98
NGC7479	32.4	39.97	109±11	2250	394	0.95	M95,TDT
NGC1068	14.4	40.38	254±10	3210	931	2.12	GM97
NGC2639	42.6	40.07	193±13	3100	401	1.67	S83
NGC3227	20.6	40.98	128±13	3168	471	0.93	NW95
NGC3982	17.0	39.15	73±4	2150	227	0.29	BHS02
NGC4051	17.0	40.25	88±13	1072	228	1.22	NW95
NGC4138	17.0	38.65	100±10	1650	211	1.07	JB96
NGC4168	16.8	38.32	187±8	2850	305	1.68	BA02
NGC4388	16.8	39.27	119±12	3900	280	0.63	TDT
NGC4565	9.7	38.42	144±7	1750	181	1.65	DB93
NGC4639	16.8	39.73	96±4	3600	180	0.32	BHS02
NGC5033	18.7	40.29	116±11	2850	261	0.85	BHS02
NGC5273	21.3	40.39	71±4	3350	147	-0.14	BHS02
NGC266	62.4	39.59		1350	326	1.99	
NGC841	59.5	39.76		1350	301	1.85	
NGC3884	91.6	40.25		2100	535	2.47	
NGC4395	3.6	38.08	30u	442	123u	0.11u	FH03
NGC4438	16.8	39.47		2050	354	1.77	
NGC4750	26.1	39.12		2200	452	2.13	
NGC4772	16.3	38.75		2400	549	2.39	
NGC5005	21.3	40.05		1650	672	3.08	
POX52	93	40.06	36±5	765	87	-0.05	BAR04
GH02	129.2	40.29	59±8	690		0.91	BG05
GH14	122.5	40.29	59±7	770		0.81	BG05
GH19	158.4	40.06	53±4	1890		-0.16	BG05

Notes:

The former 22 dwarf AGN are the objects in paper I, because, we have used new value of stellar velocity dispersion from the more recent literature for some objects, thus, we also list them in the table.

For NGC4395, the stellar velocity dispersion is the upper limit value selected from Filippenko et al. (2003). The upper limit value of [NII] is selected from Ho et al. (1997a).

References for Stellar velocity dispersion:

BA02: Bernardi et al. 2002, BAR04: Barth et al. 2004, BHS02: Barth et al. 2002, BG05: Barth, Greene & Ho 2005, DB93: DeSouza et al. 1993, F97: Fisher 1997, FH03: Filippenko & Ho 2003, GM97: Garcia-Lorenzo et al. 1997, HS98: Heraudeau & Simien 1998, JB96: Jore et al. 1996, M95: Mcelory 1995, NW95: Nelson & Whittle 1995, OF91: Ore et al. 1991, P01: Prugniel et al. 2001, PS02: Proctor & Sansom 2002, S83: Schechter 1983, S01: Sarzi et al. 2001, SP97: Simien & Prugniel 1997, SP02: Simien & Prugniel 2002, T98: Trager et al. 1998, TDT: Terlevich et al. 1990, ve01: Vega Beltran et al. 2001, z02: de Zeeuw et al. 2002

Table 2. Data of the 12 objects



Name	redshift	σ $\text{km} \cdot \text{s}^{-1}$	$L_{5100\text{\AA}}$ $10^{44} \text{erg} \cdot \text{s}^{-1}$	$\log(M_{BH})$ $M_{\odot}$	ref
Mrk590	0.02638	169 ± 28	0.653 ± 0.069	7.84 ± 0.36	nw95
3C120	0.03301	193 ± 40	1.39 ± 0.25	8.07 ± 0.48	sh90
Akn120	0.03230	177 ± 20	1.69 ± 0.12	7.92 ± 0.26	oo99
NGC3227	0.00386	128 ± 13	0.0242 ± 0.0028	7.35 ± 0.46	nw95
NGC3516	0.00884	164 ± 35	0.077 ± 0.023	7.78 ± 0.45	am97
NGC3783	0.00973	152 ± 20	0.178 ± 0.015	7.65 ± 0.27	oo95
NGC4051	0.00234	88 ± 13	0.0087 ± 0.0004	6.69 ± 0.21	nw95
NGC4151	0.00332	119 ± 26	0.0815 ± 0.0051	7.22 ± 0.41	nw95
NGC4593	0.00900	209 ± 20	0.122 ± 0.039	8.21 ± 0.25	oo99
IC4329A	0.01605	225 ± 20	0.208 ± 0.026	8.33 ± 0.24	oo99
NGC5548	0.01717	201 ± 12	0.334 ± 0.041	8.14 ± 0.17	nw95
3C390.3	0.05610	273 ± 16	0.87 ± 0.34	8.67 ± 0.22	le06

Notes:

Column V is the BH masses estimated by the equation 1a.

References for stellar velocity dispersion:

nw95: Nelson & Whittle 1995; sh90: Smith, Heckman & Illingworth 1990; oo99: Oliva et al. 1999; am97: Arribas et al. 1997; le06: Lewis & Eiraclous 2006

**Figure 1.** The correlation between the stellar velocity dispersion σ and the line width of $[\text{NII}]\lambda 6583\text{\AA}$. The solid line represents the relation $FWHM = 2.12\sigma$. The dashed line represents the relation $FWHM = 2.35\sigma$ **Figure 2.** The correlation between R_{BLRs} and $L_{H\alpha}$. The solid line represents the relation $\log(R_{BLRs}) = 0.51 \log(L_{H\alpha}/10^{44} \text{erg} \cdot \text{s}^{-1}) + 2.22$ found in paper I. The solid circles are the objects in paper I. The open circles are the ones with stellar velocity dispersions, but not included in paper I. The five-point stars are objects without stellar velocity dispersions, for which the BH masses are estimated using the line width of $[\text{NII}]\lambda 6583\text{\AA}$. The solid five-point star represents NGC 4395. The triangles are the four objects with intermediate BH masses. The objects with larger values of $L_{H\alpha}/L_{Edd}$ are marked with names or shown as triangle symbols.




Figure 3. The correlation between M_{BH} and $L_{H\alpha}$. The solid circles are the objects in paper I. The open circles are the ones with stellar velocity dispersions, but not included in paper I. The five-point stars are objects without stellar velocity dispersions, for which the BH masses are estimated using the line width of $[NII]\lambda 6583\text{\AA}$. The solid five-point star represent NGC 4395. The triangles are the four objects with intermediate BH masses. The crosses represent normal AGN.




Figure 5. The correlation between R_{BLRs} and the value of $L_{H\alpha}/L_{Edd}$. The symbols are the same as in Figure 3.




Figure 4. The correlation between M_{BH} and R_{BLRs} . The symbols are the same as in Figure 3.